

Nova V5116 Sgr and searching for superhumps in nova remnants

A. Dobrotka¹, A. Retter², A. Liu³

¹ Departement of Physics, Faculty of Materials Science and Technology, Slovak University of Technology in Bratislava, Jána Bottu 25, 91774 Trnava, The Slovak Republic, (e-mail: andrej.dobrotka@stuba.sk)

² Departement of Astronomy and Astrophysics, Penn State University, 525 Davey Lab, University Park, PA, 16802-6305, USA (e-mail: aretter@walla.com)

³ Norcape Observatory, PO Box 300, Exmouth, 6707, Australia (asliu@bigpond.net.au)

Received / Accepted

ABSTRACT

Aims. We present the period analysis of unfiltered photometric observations of V5116 Sgr (Nova Sgr 2005 #2) and we search for superhump candidates in novae remnants.

Methods. The PDM method for period analysis is used. The masses of the novae componets are estimated from the secondary mass – orbital period and primary mass – decline time relations.

Results. We found that 13 nights of V5116 Sgr observations in the year 2006 are modulated with a period of 0.1238 ± 0.0001 d (2.9712 ± 0.0024 h). Following the shape of the phased light curves and no apparent change in the value of the periodicity in different subsamples of the data, we interpret the period as orbital in nature. The binary system then falls within the period gap of the orbital period distribution of cataclysmic variables. From the maximum magnitude – rate of decline relation, we estimate the maximum absolute visual magnitude of $M_{V_{\max}} = -8.85 \pm 0.04$ mag using the measured value of decline $t_2 = 6.5 \pm 1.0$ d. The mass-period relation for cataclysmic variables yields a secondary mass estimate of about $0.26 \pm 0.05 M_{\odot}$. We propose that V5116 Sgr is a high inclination system showing an irradiation effect of the secondary star. No fully developed accretion disc up to the tidal radius with the value lower than $3.5 \cdot 10^{10}$ cm is probable. The mass ratio was estimated in a few novae and the presence or absence of superhumps in these systems was compared with the mass ratio limit for superhumps of about 0.35. We found that in the majority of novae with expected superhumps, this variability has not been found yet. Therefore, more observations of these systems is encouraged.

Key words. stars: novae, cataclysmic variables — stars: individual (V5116 Sgr) — accretion, accretion discs

1. Introduction

Novae are a subclass of cataclysmic variable stars. In these interacting binaries, the white dwarf is accreting the matter transferred from the secondary star. The accretion disc may be formed in the non magnetic case. The intermediate polar systems have a truncated disc and polar systems have magnetic field strong enough to prevent the disc formation (see Warner 1995 for review). The accumulation of critical amount of accreted material onto the white dwarf surface results in a nova explosion. The distribution of orbital periods in cataclysmic variables shows a period gap between about 2 and 3 hours. Novae do not show this lack of objects in the mentioned interval.

V5116 Sgr (Nova Sgr 2005 #2) was discovered by Liller (2005) on 2005 July 4.049 UT. The nova had a magnitude ~ 8.0 on two red photographs. An unfiltered CCD image from 2005 July 5.085 UT showed the object at mag 7.2. The spectrum from 2005 July 5.099 UT showed H_{α} with the FWHM of ~ 970 km s⁻¹. The expansion velocity derived from the sharp P Cyg profile was ~ 1300 km s⁻¹. The position of the nova was measured by Gilmore and Kilmartin (2005) and Jacques (2005). Gilmore and Kilmartin (2005) searched for the nova precursor, but no convincing candidate has been found. Russell et al. (2005) performed a $0.8 - 2.5 \mu\text{m}$ spectroscopy of the nova on 2005 July 15. The

object showed emission lines of H I, He I, C I, N I, Ca II and O I with a FWHM ~ 2200 km s⁻¹. He I showed P Cyg profile at 1.0830 and 2.0581 μm . No thermal dust emission was observed.

After the nova explosion the accretions disc is destroyed. The new disc is reformed by the stream of matter flowing from the secondary, interacting with itself and forming a ring in the circularisation radius. The disc starts to form by the viscous shearing. Matter losing angular momentum is moving invard and the excess of angular momentum is transported by the matter flowing outward. The invard moving matter touches the white dwarf and the disc is reformed (Pringle 1981). If the white dwarf is magnetic, the matter interacts with the magnetosphere and is then conducted by the magnetic field to the magnetic poles. The interaction of the flow with the poles of the rotating star is observed as periodic signal with the spin period of the white dwarf. In the case of intermediate polars the spin period is usually much shorter than the orbital period (Patterson 1994, Hellier 1996). The polar systems without a disc are synchronous rotators, hence the spin period is equal or close to the orbital period (see e.g. Schmidt and Stockman 1991). Nova V1500 Cyg (polar system without a disc) changed the period from 0.141 d to 0.138 d and then stabilised at 0.140 d (Patterson 1978, 1979). The difference of 1.8% between the rotation period of the white dwarf and the orbital period is ascribed to the effects of the nova explosion in 1975.

The synchronisation can be then corrupted by the nova explosion, but the spin period remain still very close to the orbital period. This is in contrast with intermediate polars with disc presence.

Searching for periods in novae allows to study the orbital distribution and evolution of these systems. Currently, there are about 50 novae with known orbital periods (Warner 2002) with typical values ranging from 2 to 9 hours. The existence of the accretion disc or its renovation after the nova explosion is confirmed by the detection of the superhump period (see e.g. Retter et al. 1997, Kang et al. 2006a) or the spin period of a magnetic white dwarf in the case of intermediate polars (see e.g. Retter et al. 1998).

Superhumps are caused by a precessing accretion disc generally in systems with mass ratio $M_2/M_1 < 0.35 \pm 0.02$ (see Patterson et al. 2005 for review), in which the disc radius reaches the 3:1 resonance. The mass ratio indicates that systems with massive primaries and low mass secondaries are probable superhumpers. Novae in general possess high mass white dwarfs (see e.g. Warner 1995, Smith and Dhillon 1998, Webbink 1990) and short orbital periods suggest low mass secondaries. It is therefore meaningful to search for superhump variability in novae with short orbital period.

We have an ongoing program to observe novae with small telescopes to search for periodicities in their optical light curves. In this paper we report the detection of one periodicity in our photometric data ($P = 0.1238 \pm 0.0001$ d) of V5116 Sgr and we discuss the presence of superhumps in nova remnants. In Section 2 we present our observational material. The long-term light curve with the period analysis of the data is presented in Sec. 3 and in Sec. 4 we discuss the long-term light curve behaviour (Sec. 4.1), the results of the period analysis (Sec. 4.2) and superhump search in nova remnants (Sec. 4.3).

2. Observations

V5116 Sgr was observed during 13 nights in 2006. The observations included 45.2 hours and 1256 data points in total. The list of observations is presented in Table 1. The photometry was carried out with a 0.3-m f/6.3 telescope coupled to a SBIG ST7E CCD camera. The telescope is located in Exmouth, Western Australia. The pixel size of the CCD is 9×9 microns. This camera is attached to an Optec f5 focal reducer giving an image field of view of 15×10 arcmin. The range of seeing for the data was 2.5 – 3 arcsec. The exposure times were between 30 and 90 sec every 120 sec, and no filter was used. Aperture photometry was used in the reduction, with an aperture size of 12 pixels (radius). As comparison star we used GSC 7392 856 with $V = 12$ mag and as check star GSC 7392 292 with $V = 11.6$ mag. The instrumental magnitudes of the stars were derived from the SBIG CCDOPS software and the magnitudes of the comparison and check stars from the GSC.

Figure 1 displays our light curve and the long-term light curve of the nova from the American Association of Variable Star Observers (AAVSO). In Fig. 2 we show 4 best examples of our observational runs with the sinusoidal fit (with first harmonic) to the data using the period derived in this paper. The studied variations are clearly seen.

Table 1. The observational log. HJD is -2453000 days, t is the duration of the observation in hours and N is the number of frames.

Run	Date	HJD (start)	t [h]	N
1	2006 August 30	977.9654	4.8	133
2	2006 August 31	978.9766	4.5	123
3	2006 September 21	999.9753	3.4	94
4	2006 September 22	1000.9831	3.3	93
5	2006 September 24	1002.9704	3.6	101
6	2006 September 27	1005.9734	3.5	93
7	2006 September 28	1006.9792	3.3	90
8	2006 September 29	1007.977	3.4	98
9	2006 October 04	1012.974	3.4	94
10	2006 October 08	1016.9812	3.2	90
11	2006 October 10	1018.9811	3.0	82
12	2006 October 15	1023.9679	3.2	91
13	2006 October 18	1026.9677	2.6	74




Fig. 1. The AAVSO light curve (upper panel) and the unfiltered light curve of our data (bottom panel).

3. Data analysis

3.1. Long-term light curve

By using the maximum visual magnitude of $V = 7.15$ we measured the time required for a decline of two magnitudes from maximum $t_2 = 6.5 \pm 1.0$ d and three magnitudes from maximum $t_3 = 20.2 \pm 1.9$ d (Fig. 1 – upper panel). This makes V5116 Sgr a very fast nova according to the classification given in Table 5.4 of Warner (1995). We calculated the visual absolute magnitude at maximum brightness using Equation (2) of Della Valle and Livio (1995) and obtained $M_{V\max} = -8.85 \pm 0.04$ mag.

The bottom panel of the Fig. 1 shows our unfiltered photometry of the nova. The long-term variations with time scale $\sim 30 - 40$ days with amplitude ~ 1 mag ~ 450 days after the outburst is clear. Similar large scale quasi-periodic




Fig. 2. A selection of our observations (4 detrended best runs; 1,2,11 and 13). The solid curve is the sinusoidal fit (plus first harmonic) to the data using the period derived in this paper.

brightness oscillations with amplitude 1 – 1.5 mag months after the maximum were observed for example in GK Per, V603 aql, DK Lac (see e.g. Warner 1995) or V4745 Sgr (Csák et al. 2005), with time scales 5,12,25 and ~ 20 days respectively.

3.2. Period analysis

For the period analysis we used the PDM method (Stellingwerf 1978). First, the data were detrended by subtracting the linear fit from each night. A simple visual inspection of the runs suggests that the data are modulated with a periodicity of ~ 1.5 hours. This value was then identified as the first harmonic of the real periodicity, therefore the exact value of the frequency is $f = 8.080 \pm 0.006 \text{ d}^{-1}$, which corresponds to the periodicity $P = 0.1238 \pm 0.0001 \text{ d}$ ($2.9712 \pm 0.0024 \text{ h}$). We calculated the errors in the frequency from the half width at the half minimum of the peaks in the PDM power spectrum.

The results from the period analysis of all the data are depicted in Fig. 3. The suspected periodicity ($f = 8.080 \pm 0.006 \text{ d}^{-1}$, $P = 0.1238 \pm 0.0001 \text{ d}$) is seen with its 1-d aliases and the dominant peak is the first harmonic ($2 \times f = 16.160 \pm 0.006 \text{ d}^{-1}$). For testing which value is the real signal and which is the alias we performed an O–C diagram with a linear fit to the data (Fig. 4; upper panel – O–C for the frequency 8.08 d^{-1} and lower panel – O–C for the frequency 9.08 d^{-1}). The lower panel shows a larger scatter of the O–C values compared with the upper panel. This indicates that the real signal is $f = 8.080 \pm 0.006 \text{ d}^{-1}$ and the dominant peak at $\sim 16 \text{ d}^{-1}$ is the first harmonic of the real signal. To test the reality of the peaks we performed a power spectrum of the synthetic data using a sinusoids with the derived periods (including fundamental f , first




Fig. 3. Power spectra of all the data. Panel a – raw detrended data (night means subtracted), panel b – data after the subtraction of the harmonic frequency $2 \times f$, panel c – data after the subtraction of the harmonic $2 \times f$ and the fundamental frequency f , panel d – data after the subtraction of the harmonic $2 \times f$, the fundamental frequency f and the harmonic frequency $4 \times f$, panel e – synthetic spectrum with the frequencies f , $2 \times f$ and $4 \times f$. The dashed line represent $f = 8.08 \text{ d}^{-1}$.

harmonic $2 \times f$ and $4 \times f$ described later) and phases with amplitudes found from the fits to the data sampled as the original data with introduced Gaussian noise (Fig. 3e). The similarity with the original periodogram is clear. Further we performed a period analysis of two different subsamples of the data (first subsample – runs 3 – 8; second subsample – runs 9 – 13) depicted in Fig. 5. The frequency is present together with its first harmonic in both datasets. The values are $8.090 \pm 0.015 \text{ d}^{-1}$ and $8.080 \pm 0.028 \text{ d}^{-1}$ (first and second subsample respectively) which is within the errors in both cases. We also performed an analysis of the check star and did not identify any periodicity associated with its data.

There is a strong peak near the value $\sim 8.58 \text{ d}^{-1}$. This signal is not real as seen in Fig. 3b,c. After the subtraction of $2 \times f$ (Fig. 3b) this peak disappeared and after a further subtraction of f (Fig. 3c) no peak is dominant in this frequency region. This feature also appears in the synthetic power spectrum (Fig. 3e), so it is definitely an alias of f .

Another strong feature is a peak at $\sim 35 \text{ d}^{-1}$. Subtracting the harmonic frequency $4 \times f$ from the data, after the subtraction described in the previous paragraph, this signal disappeared (Fig. 3d). It is another artefact of the real signal f . No other dominant peaks are visible with frequencies higher than 40 d^{-1} . No other frequency was then found.




Fig. 4. O-C diagram. Upper panel – for frequency $8.080 \pm 0.006 \text{ d}^{-1}$ ($P = 0.1238 \pm 0.0001 \text{ d}$) and lower panel – for frequency $9.080 \pm 0.006 \text{ d}^{-1}$ ($P = 0.1101 \pm 0.0001 \text{ d}$). The solid line is the linear fit to the data.




Fig. 5. Power spectra of two different subsamples of the data. Upper panel – runs 3 – 8; lower panel – runs 9 – 13. The dashed line represents $f = 8.08 \text{ d}^{-1}$.

3.3. The structure of the periodicity

In Figure 6 we show the light curve of V5116 Sgr folded on the 0.1238 d period. The points are averaged magnitudes in 50 equal bins that cover the 0 – 1 phase interval. The full amplitude of the mean variation is $0.43 \pm 0.02 \text{ mag}$ for primary minimum and $0.30 \pm 0.02 \text{ mag}$ for secondary




Fig. 6. The light curve of V5116 Sgr folded on the 0.1238 d period and binned into 50 equal bins. Panel a – all the data, panel b – runs 3 – 8; panel c – runs 9 – 13.

minimum. Using the sinusoidal fit to determine the amplitude is not adequate, because of the non sinusoidal shape of the light curve, we therefore estimated the amplitude by measuring the difference between extrema. The bars are the errors of the mean value. The light curves folded on the periodicity found show typical features of an eclipsing system.

We determined the epoch of minima by fitting low-order polynomials to selected parts of the light curve. The best fitted (using sinusoidal fits) ephemeris of the periodicity is:

$$T_{(\min)}(HJD) = 2453978.0726 + 0.1238E \\ \pm 0.0009 \pm 0.0001$$

4. Discussion

4.1. Long-term variations of V5116 Sgr

The long-term light curve of the nova V5116 Sgr (Fig. 1 – bottom panel) shows a transition from smooth decline to probable oscillations. Several models have been suggested for such “transient” phase. One of them explains the transition as the time when the accretion disc is re-established (Retter 2002). The authors propose a connection between this phase and intermediate polars. There is no indication of magnetic nature of the white dwarf in V5116 Sgr yet. Spin period of the white dwarf indicating the intermediate polar type of this “transient” phase in novae was detected for example in V4745 Sgr by optical observations (Dobrotka et al. 2006a), V1494 Aql (Drake et al. 2003) and V4743 Sgr by X-ray observations (Ness et al. 2003). If this “transient” phase interpretation of Retter (2002) is applicable in the case of V5116 Sgr, the accretion must be then re-established and the accretion disc should be present.

4.2. Short-term variations of V5116 Sgr

We have identified one periodicity in the light curve of the nova V5116 Sgr about 15 months after the maximum brightness. The value is $P = 0.1238 \pm 0.0001$ d (2.9712 ± 0.0024 h). The upper limits of the period difference between two subset of data with $\simeq 16$ days of mean distance analysed in Fig. 5 is $1.5 \cdot 10^{-4}$ d which gives $|\dot{P}| \simeq 0.94 \cdot 10^{-5}$. Following Patterson et al. (1993) the mean variation of the superhump period in SU UMa superhumping systems is $|\dot{P}| \simeq 3 - 10 \cdot 10^{-5}$ (see their Table. 1). For the recurrent nova VY Aqr the authors derived a variation of $|\dot{P}| \simeq 8.2 \cdot 10^{-5}$. Our value of $|\dot{P}|$ comes probably from the errors of period measurements rather than from real period changes. The period seems then to be constant during our observations and the shape of the folded light curve suggests a primary and a secondary eclipse. We therefore propose that this periodicity is the orbital period of the binary system. Such a period is at the lower end of the mostly populated region of orbital periods in novae (Warner 2002). The dominant first harmonic frequency in the power spectra is a result of the clear structure primary – secondary eclipse in the folded light curve which suggests a high inclination angle of the binary system. Using the orbital period of the system and equation (9) from Smith and Dhillon (1998) we obtain a rough estimate for the secondary star mass of $0.26 \pm 0.05 M_{\odot}$. Using a mean white dwarf mass of $0.85 \pm 0.05 M_{\odot}$ from Smith and Dhillon (1998), we find a mass ratio (secondary/primary mass) of $M_2/M_1 = 0.3 \pm 0.1$.

After the nova explosion, the hot white dwarf may heat and irradiate the cooler companion. The observed orbital light curve of the nova can be the result of the aspect variations or eclipses of the secondary due to heating from the hot primary and the asymmetry in the pulse profiles could be produced once the shape of the secondary is of a tear drop model. The irradiation effects in classical novae can also be detected long after the outburst stage (e.g., V1500 Cyg; Sommers and Naylor 1999, DN Gem; Retter et al. 1999). Two eclipse like features are present in the folded light curve of V5116 Sgr (Fig. 6). The shape is similar to the light curve of V2540 Oph in 2003 (Ak et al. 2005) with a dip at phase ~ 0.5 ($\sim 33\%$ amplitude of the primary minimum for V2540 Oph and $\sim 70\%$ for V5116 Sgr). The authors concluded that V2540 Oph is likely a high inclination system either showing an irradiation effect or having a spiral structure in its accretion disc. Woudt and Warner (2003a) noted that one of the following requirements must be fulfilled for the large amplitude orbital modulations to be seen in the light curve of a recent nova in which the accretion disc does not dominate the luminosity of the system: 1) the disc is foreshortened but the secondary is seen (a high inclination angle), 2) the disc has small dimensions, 3) no disc (case of polars). In the case of V5116 Sgr the polar interpretation is possible because of the orbital period distribution of polars which peaks below 5 hours (Warner 1995), but nothing else indicate this option. The small dimension of the disc is supported by; 1) the short orbital period, 2) the post nova stage when the disc is reforming after explosion.

A comparison to two novae within the period gap IM Nor and DD Cir with present irradiation effect (Woudt and Warner 2003b,c) can be made. The light curve of V5116 Sgr is different from those of IM Nor and DD Cir but similar to V2540 Oph. IM Nor and DD Cir showed

very small dips at phase 0.5 interpreted as partial eclipses of the irradiated secondary by the disc or matter stream. The light curve shape depends on the disc radius and on the inclination angle. The deep secondary eclipses in V5116 Sgr require a large disc or a high inclination. The strength of the irradiation effect (white dwarf post-outburst temperature) can also play a significant role. The differences in the phase of the maxima in Fig. 6b,c can be due to spiral structures in the disc as mentioned before in regard with the nova V2540 Oph.

The reconstruction of an accretion disc after the nova eruption is indicated by the discovery of the white dwarf spin or by the superhump period. A probable spin period of the white dwarf was detected ~ 1 year after the outburst in V4745 Sgr (Dobrotka et al. 2006a), ~ 2.75 years in V4743 Sgr (Kang et al. 2006b) and ~ 15 months after the maximum in V1425 Aql (Retter et al. 1998). Several systems show superhumps as early as two and a half months after the outburst like V4633 Sgr (a spin period is another option in this case, Lipkin et al. 2001) or two years after the outburst like V1974 Cyg (Retter et al. 1997). In our V5116 Sgr light curves extending ~ 470 days after outburst we did not find any photometric variations consistent with white dwarf spin modulation or superhump properties. We can not say anything about the magnetic nature of the white dwarf, but the components mass ratio indicates that the superhump existence is probable. The mass ratio using the primary mass average is not decisive. Using the derived secondary mass $0.26 M_{\odot}$ (Section 4.2) the mass ratio is lower than 0.35 for primary mass higher than $0.74 M_{\odot}$. Taking all known primary masses in cataclysmic variables (Ritter and Kolb 2003), 63% have higher or equal mass than $0.74 M_{\odot}$. This probability value is not enough to make sure that superhumps are expected in V5116 Sgr, but the search for such variability could be fruitful. However V5116 Sgr is a very fast nova. According to equation (13) from Livio (1992) and taking $t_3 = 20.2 \pm 1.9$ d derived in this paper (Section 3.1) we obtain a mass estimate of the white dwarf of $1.04 \pm 0.02 M_{\odot}$ and thus $M_2/M_1 = 0.25 \pm 0.05$. The presence of superhumps is then expected if a disc is fully developed up to tidal radius.

The superhump period is a few percent longer or shorter than the orbital period. The possibility that the periodicity found in this paper is a superhump is rejected following the stability discussion and the shape of the folded light curve (Fig. 6). The mean shape of superhumps is typically an asymmetric sinusoid and our data show typical eclipse like features.

The presence of the disc in cataclysmic variables depends on the mass loss from the secondary. The material supplied from the secondary within the period gap depends on the strength of the magnetic braking driving the secondary out of thermal equilibrium (mass loss time scale shorter than the thermal time scale). The stronger the braking, the wider the gap will be and the higher is the upper end of the period gap. If the magnetic braking is low enough, the mass loss time scale may never become shorter than the thermal time scale. In the case of novae another condition is important. The nova explosion heats the secondary which leads to an enhanced mass transfer. Therefore the complete absence of accretion discs in novae within the period gap is not expected.

4.3. Searching for superhumps in other nova remnants

The orbital period 2.462 h of the mentioned IM Nor (Woudt and Warner 2003b) yields a secondary mass estimate of $0.20 M_{\odot}$ following Smith and Dhillon (1998). The decay time $t_3 \simeq 50$ d (Kato et al. 2002) yields a primary mass estimate of $\simeq 0.86 M_{\odot}$ (Livio 1992) giving a mass ratio $\simeq 0.23$. Woudt and Warner (2003b) concluded that they did not observe the superhump modulation. This conclusion with the mass ratio safely lower than 0.35 indicate that the existence of a fully developed disc is not probable. DD Cir (Woudt and Warner 2003c) has a period of 2.339 h yielding a secondary mass estimate of $0.18 M_{\odot}$. The decay time $t_2 \simeq 4.5$ d (Liller 1999, no other estimates are available) place the object in the class of very fast novae (similar to V5116 Sgr). The decay time $t_3 \simeq 10$ d following the equation $t_3 \simeq 2.75 t_2^{0.88}$ from Warner (1995) gives a rough white dwarf mass estimate of $\simeq 1.16 M_{\odot}$ (mass ratio < 0.2). Woudt and Warner (2003b) interpreted the deep eclipse as obscuration of the disc and did not detect any periodicity near the orbital period in the period analysis. Therefore, following the tidal instability model it is again probable that the disc is not fully developed up to 3:1 resonance radius 3 years after the outburst. However the authors argued that the disc radius is 47 % of the orbital separation. Using the third Kepler law we obtain a disc radius of $\sim 3.1 \cdot 10^{10}$ cm. The disc in DD Cir is then large enough to reach the 3:1 resonance radius calculated from the equation (3.39) from Warner (1995); $r_{3:1} \sim 3 \cdot 10^{10}$ cm. The result is marginally in contrast with the absence of superhumps. Taking the derived values for V5116 Sgr we get $r_{3:1} \simeq 3.5 \cdot 10^{10}$ cm as an upper limit of the disc radius taking the absence of superhumps into consideration. The boundaries of the eclipse in our light curve are not so clear as in the DD Cir case, therefore not suitable to estimate the disc radius.

Comparison to other systems follows the same way as in the case of IM Nor and DD Cir. We took components masses from the Catalogue of Ritter and Kolb (2003) and other masses are estimated from the orbital period (using Smith and Dhillon 1998) and t_3 time (using Livio 1992). We rejected systems with unknown or non measurable t_2 or t_3 time (peculiar light curve, strong oscillations in the decline) without catalogue mass values (no information on mass ratio, ex: RS Car), magnetic novae (AM Her systems without disc, ex: V1500 Cyg), systems with orbital period > 10 h (Smith and Dhillon fitting not adequate, systems with evolved secondary are then excluded too, ex: DI Lac, V841 Oph, V723 Cas, GK Per), systems with insufficient photometric observations to detailed period analysis (ex: V500 Aql, V446 Her, HZ Pup, DY Pup, CT Ser). The final list of systems is in Table 2.

The results are depicted in Fig. 7. The lower panel only shows the systems with known catalogue mass values and the upper panel is after the addition of estimated mass values in this work. Filled circles are systems with detected superhumps. The critical mass ratio of 0.35 is also shown. The results presented in the lower panel are as expected from the mass ratio limit for superhumps, but the statistical set is small. Adding other mass estimates changes the situation. 7 systems below or close to the limit 0.35 show superhumps but 14 systems do not. The estimated mass ratio has a mean difference of 0.2 from the catalogue values. The differences are distributed randomly. Therefore there is




Fig. 7. Selected nova remnants in the mass ratio (q) – orbital period (P_{orb}) plane. Open circle – non detected superhumps, filled circle – detected superhumps. Panel a – mass ratio from combined sources (catalogue and estimates from P_{orb} , t_3 time), panel b – mass ratio only from catalogue values (Ritter and Kolb 2003). The dashed line is the 0.35 limit.

no reason to suspect that all mass ratios are systematically underestimated. The 7 systems with detected superhumps are safely located in comparing to the 0.35 critical value as expected by theory and are only a third of all systems falling below or close to the mass ratio of 0.35. The majority of all systems occupy the orbital period interval of 2 – 4 hours. Following Patterson (2005) this interval has a $\sim 40 - 90$ % probability to observe superhumps. 7 systems versus 14 from our investigation yields only ~ 30 %.

Superhumps are observed during the outbursts of short orbital period dwarf novae (SU UMa stars) where the disc reaches the 3:1 resonance radius (superoutburst case). The disc during this active stage (outburst – active and hot stage, quiescence – not active and cold stage) reaches larger radius than in quiescence (see Lasota 2001 for review). The irradiation of the secondary by the central white dwarf in nova remnants can produce enhanced mass transfer rate strong enough to keep the disc in this stable hot active stage (as in the nova like stars) with the larger diameter (Hameury and Lasota 2005). In addition, superoutbursts in SU UMa stars, when superhumps are observed, can be explained by the irradiation and the enhanced mass transfer (Smak 1995, 2004, Hameury et al. 2000, Schreiber et al. 2004). Therefore, permanent superhumpers are possible after a nova outburst. A few possible interpretation of the superhump lack can be: 1) not systematically applicable tidal instability theory; tidal torques are probably not the main and only condition of superhump existence as concluded by Hameury and Lasota (2005), 2) disc radius not developed up to tidal radius, 3) superhumps present but not detected, because of insufficient sets of data.

5. Summary

By using the maximum visual magnitude of $V = 7.15$ we measured the decay time $t_2 = 6.5 \pm 1.0$ d from the long-term light curve. This makes V5116 Sgr a very fast nova. We estimated a visual absolute magnitude in maximum of $M_{V\max} = -8.85 \pm 0.04$ mag. We found a period of 0.1238 ± 0.0001 d (2.9712 ± 0.0024 h) which we interpret as the orbital period of the binary system. We propose that V5116 Sgr is a high inclination system showing a strong irradiation effect of the secondary star. No fully developed accretion disc up to the tidal radius is possible with radius lower than $3.5 \cdot 10^{10}$ cm.

Searching for superhumps in nova remnants using the limiting value for mass ratio 0.35 and estimated mass values shows, that in the majority of systems with expected superhumps this variability has not been found yet. Looking for superhumps in these systems could be fruitful.

6. Acknowledgments

This work was supported by the Slovak Academy of Sciences Grant No. 2/7011/7. We acknowledge with thanks the variable star observations from the AAVSO International Database contributed by observers worldwide and used in this research. We also acknowledge Michael Friedjung for valuable comments.

References

- Ak, T., Retter, A., Liu, A., 2005, *PASA* 22, 298
- Arhipova, V.P., Burlak, M.A., Esipov, V.F., 2002, *Astronomy Letters* 28, 100
- Balman, S., Yilmaz, A., Retter, A., Saygac, T., Esenoglu, H., 2005, *MNRAS* 356, 773
- Balman, S., Retter, A., Bos, M., 2006, *AJ* 131, 2628
- Baptista, R., Jablonski, F.J., Cieslinski, D., Steiner, J.E., 1993, *ApJ* 406, L67
- Barlow, M.J., Brodie, J.P., Brunt, C.C., Hanes, D.A., Hill, P.W., Mayo, S.K., Pringle, J.E., Ward, M.J., Watson, M.G., Whelan, J.A.J., Willis, A.J., 1981, *MNRAS* 195, 61
- Barsukova, E.A., Goranskii, V.P., 2003, *AstL* 29, 195
- Csák, B., Kiss, L.L., Retter, A., Jacob, A., Kaspi, S., 2005, *A&A* 429, 599
- Campbell, R.D., Shafter, A.W., 1995, *ApJ* 440, 336
- Della Valle, M., Livio, M., 1995, *ApJ* 452, 704
- Della Valle, M., Pasquini, L., Daou, D., Williams, R.E., 2002, *A&A* 390, 155
- Diaz, M.P., Bruch, A., 1994, *IBVS* 4079, 1
- Diaz, M.P., Bruch, A., 1997, *A&A* 322, 807
- Dobrotka, A., Retter, A., Liu, A., 2006a, *MNRAS* 371, 459
- Dobrotka, A., Friedjung, M., Retter, A., Hric, L., Novák, R., 2006b, *A&A* 448, 1107
- Drake, J.J., Wagner, R.M., Starrfield, S., Butt, Y., Krautter, J., et al., 2003, *ApJ* 584, 448
- Duerbeck, H.W., 1987, *Sp. Sci. Rev.* 45, 1
- Gilmore, A.C., Kilmartin, P.M., 2005, *IAUC* 8559
- Hameury, J.-M., Lasota, J.-P., Warner, B., 2000, *A&A* 353, 244
- Hameury, J.-M., Lasota, J.-P., 2005, *A&A* 443, 283
- Hellier, C., 1996, in “Cataclysmic Variable and Related Objects”, eds. Evans, N., Wood, J.H., *IAU Colloq* 158, Kluwer, Dordrecht, 143
- Hoffmeister, C., 1959, *AN* 285, 149
- Jacques, C., 2005, *IAUC* 8559
- Kaluzny, J., 1990, *MNRAS* 245, 547
- Kang, T. W., Retter, A., Liu, A., Richards, M. 2006a, *AJ* 131, 1687
- Kang, T. W., Retter, A., Liu, A., Richards, M. 2006b, *AJ* 132, 608
- Kato, T., Yamaoka, H., Liller, W., Monard, B., 2002a, *A&A* 391, 7
- Kato, T., Hishikura, T., West, J.D., et al., 2002b, *IAUC* 7976
- Kato, T., Ishioka, R., Uemura, M., Starkey, D., Krajci, T., 2004, *PASJ* 56, 125
- Kiss, L.L., Gogh, N., Vinkó, et al., 2002, *A&A* 384, 982
- Kohoutek, L., Pauls, R. 1980, *A&A* 92, 200
- Lasota, J.-P., 2001, *New Asron. Rev.* 45, 449
- Leibowitz, E.M., Mendelson, H., Mashal, E., 1992, *ApJ* 385, L49
- Lipkin, Y., Leibowitz, E.M., Retter, A., Shemmer, O., 2001, *MNRAS* 328, 1169
- Liller, W., 2005, *IAUC* 8559
- Livio, M., 1992, *ApJ* 393, 516
- Lynch, D.K., Wilson, J.C., Rudy, R.J., Venturini, C., Mazuk, S., Miller, N.A., Puetter, R.C., 2004, *AJ* 127, 1089
- Matsumoto, K., Uemura, M., Kato, T., Kiyota, S., Ayani, K., Kawabata, T., Král, L., Havlík, T., Kolasa, M., Novák, R., Masi, G., 2001, *A&A* 378, 487
- Ness, J.U., Starrfield, S., Burwitz, V., Wichmann, R., Hauschildt, P., et al., 2003, *ApJ* 594, L127
- Paczynski, B., 1971, *Acta Astronomica* 21, 417
- Patterson, J., 1978, *ApJ* 225, 954
- Patterson, J., 1979, *ApJ* 231, 789
- Patterson, J., 1979a, *ApJ* 233, L13
- Patterson, J., Bond, H.E., Grauer, A.D., Shafter, A.W., Mattei, J.A., 1993, *PASP* 105, 69
- Patterson, J., 1994, *PASP* 106, 697
- Patterson, J., Warner, B., 1998, *PASP* 110, 1026
- Patterson e al., 2005, *PASP* 117, 1204
- Pringle, J.E., 1981, *Ann.Rev.Astron.* 19, 137
- Retter, A., Leibowitz, E.M., Ofek, E.O., 1997, *MNRAS* 286, 745
- Retter, A., Leibowitz, E.M., 1998, *MNRAS* 296, L37
- Retter, A., Leibowitz, E.M., Kovo-Kariti, O., 1998, *MNRAS* 293, 145
- Retter, A., Leibowitz, E.M., Naylor, T., 1999, *MNRAS* 308, 140
- Retter, A., Naylor, T., 2000, *MNRAS* 319, 510
- Retter, A., Lipkin, Y., 2001, *A&A* 365, 508
- Retter, A., 2002, in “Classical Nova Explosion”, eds. Hernanz, M., José, J., *AIP Conf. Ser.* 637, 279
- Ritter, H., Kolb, U., 2003, *A&A* 404, 301
- Robinson, E.L., Nather, R.E., Kepler, S.O., 1982, *ApJ* 254, 646
- Russell, R.W., Lynch, D.K., Rudy, R.J., 2005, *IAUC* 8579
- Schmidt, G.D., Stockman, H.S., 1991, *ApJ* 371, 749
- Schreiber, M.R., Hameury, J.-M., Lasota, J.-P., 2004, *A&A* 427, 621
- Shafter, A.W., Misselt, K.A., veal, J.M., 1993, *PASP* 105, 853
- Shafter, A.W., Misselt, K.A., 1995, *ApJ* 448, L33
- Skillman D.R., Harvey, D., Patterson, J., Vanmunster, T., 1997, *PASP* 109, 114
- Sommers, M.W., Naylor, T., 1999, *A&A* 352, 563
- Smak, J., 1995, *AcA* 45, 355
- Smak, J., 2004, *AcA* 54, 221
- Smith, D.A., Dillon, V.S., 1998, *MNRAS* 301, 767
- Stellingwerf, R.F., 1978, *ApJ* 224, 953
- Thorstensen, J.R., Cynthia, J.T., 2000, *MNRAS* 312, 629
- Vogt, N., 1975, *A&A* 41, 15
- Walker, M.F., 1961, *ApJ* 134, 171
- Walker, M.F., 1963, *ApJ* 138, 313
- Warner, B., 1986, *MNRAS* 219, 751
- Warner, B., 1986, *MNRAS* 222, 11
- Warner, B., 1995, *Cataclysmic Variable Stars*, Cambridge University Press, Cambridge
- Warner, B., 2002, in “Classical Nova Explosions”, eds. Hernanz, M., José, J., *AIP Conf. Ser.* 637, 3
- Webbink, R.F., 1990, in “Accretion-Powered Compact Binaries”, ed. Mauche, C.W., Cambridge University Press, Cambridge, 177
- Woudt, P.A., Warner, B., 2001, *MNRAS* 328, 159
- Woudt, P.A., Warner, B., 2002, *MNRAS* 335, 44
- Woudt, P.A., Warner, B., 2003a, *MNRAS* 339, 731
- Woudt, P.A., Warner, B., 2003b, *MNRAS* 343, 313
- Woudt, P.A., Warner, B., 2003c, *MNRAS* 340, 1011
- Zhao, P., McClintock, 1997, *ApJ* 483, 899

Table 2. Selected nova systems. P_{orb} is the orbital period in hours, M_2 is the secondary mass in solar masses derived from Smith and Dhillon (1998) (from Ritter and Kolb 2003 in parentheses), t_3 is the decline time of three magnitudes from maximum (value in parentheses is the value estimated from the t_2 time using $t_3 \simeq 2.75 t_2^{0.88}$ from Warner 1995), M_1 is the primary mass derived from Livio (1992) (from Ritter and Kolb 2003 in parentheses), q is the mass ratio, SH is the information about superhump detection (Y – yes, N – no) and Ref. is the reference.

System	P_{orb} [h]	M_2 [M_{\odot}]	t_3 [d]	M_1 [M_{\odot}]	q	SH	Ref.
RW UMi	1.418	0.07	140	0.64	0.11	N	19
CP Pup	1.474	0.08	8	1.19	0.07	Y	22
V1974Cyg	1.950	0.14	42	0.90	0.16	Y	9,26
RS Car	1.980	0.14	(58)	0.83	0.17	N	24
DD Cir	2.339	0.18	(10)	1.16	0.16	N	4
IM Nor	2.462	0.20	50	0.86	0.23	N	3,13
QU Vul	2.682	0.23	40	0.91	0.25	N	29
V2214 Oph	2.804	0.24	100	0.71	0.34	N	17
V630 Sgr	2.830	0.25	11	1.14	0.22	Y	23
V351 Pup	2.837	0.25	(32)	0.95	0.26	N	23
V5116 Sgr	2.971	0.26	20.2	1.04	0.25	N	this paper
V4633 Sgr	3.014	0.27	42	0.90	0.30	Y	14
DN Gem	3.068	0.28	37	0.92	0.30	N	28
V1494 Aql	3.231	0.30	16	1.08	0.28	N	8,30,31
V1668 Cyg	3.322	0.31	23	1.02	0.30	N	32
V603 Aql	3.324	0.31(0.29)	8	1.19(1.2)	0.26(0.24)	Y	21
RR Cha	3.370	0.31	58	0.83	0.37	Y	24,1
RR Pic	3.481	0.33(0.4)	150	0.62(0.95)	0.53(0.42)	N	33,34
V382 Vel	3.508	0.33	9	1.17	0.28	N	2,12
V533 Her	3.530	0.33	44	0.89	0.37	N	35,42
V2574 Oph	3.550	0.34	(33)	0.95	0.36	Y	15
OY Ara	3.731	0.36(0.34)	...	(0.82)	(0.41)	N	36
V1493 Aql	3.740	0.36	7	1.21	0.30	N	8,11
V849 Oph	4.146	0.41	175	0.59	0.69	N	37
DQ Her	4.647	0.48(0.40)	94	0.72(0.60)	0.67(0.67)	N	38
T Aur	4.906	0.51(0.63)	100	0.71(0.68)	0.72(0.92)	N	41
V4745 Sgr	4.987	0.52	32.8	0.95	0.55	N	7,10
HR Del	5.140	0.54(0.55)	230	0.54(0.67)	1.00(0.82)	N	43
V1425 Aql	6.139	0.57	39	0.91	0.63	N	8,27
V4743 Sgr	6.718	0.74	15	1.10	0.67	N	16,39
BY Cir	6.760	0.74	157	0.62	1.19	N	4
V2540 Oph	6.835	0.75	213	0.55	1.36	N	18
V838 Her	7.143	0.79(0.74)	3.2	1.29(0.87)	0.61(0.85)	N	44
V2275 Cyg	7.548	0.84	7	1.21	0.69	N	20,40
BT Mon	8.012	0.90(0.87)	(213)	0.55(1.04)	1.64(0.84)	N	45
V368 Aql	8.285	0.93	42	0.90	1.03	N	46
QZ Aur	8.580	0.97(1.05)	27	0.99(1.05)	0.98(1.00)	N	47

References:

1 - Hoffmeister (1959), 2 - Balman et al. (2006), 3 - Woudt and Warner (2003b), 4 - Woudt and Warner (2003c), 5 - Lynch et al. (2004), 6 - Barlow et al. (1981), 7 - Csák et al. (2005), 8 - Arhipova et al. (2002), 9 - Skillman et al. (1997), 10 - Dobrotka et al. (2006a), 11 - Dobrotka et al. (2006b), 12 - Della Valle et al. (2002), 13 - Kato et al. (2002a), 14 - Lipkin et al. (2001), 15 - Kang et al. (2006a), 16 - Kang et al. (2006b), 17 - Baptista et al. (1993), 18 - Ak et al. (2005), 19 - Retter and Lipkin (2001), 20 - Balman et al. (2005), 21 - Retter and Naylor (2000), 22 - Patterson and Warner (1998), 23 - Woudt and Warner (2001), 24 - Woudt and Warner (2002), 25 - Matsumoto et al. (2001), 26 - Retter et al. (1997), 27 - Retter and Leibowitz (1998), 28 - Retter et al. (1999), 29 - Shafter and Misselt (1995), 30 - Kato et al. (2004), 31 - Barsukova and Goranskii (2003), 32 - Kaluzny (1990), 33 - Vogt (1975), 34 - Warner (1986), 35 - Patterson (1979a), 36 - Zhao and McClintock (1997), 37 - Shafter et al. (1993), 38 - Walker (1961), 39 - Kato et al. (2002b), 40 - Kiss et al. (2002), 41 - Walker (1963), 42 - Thorstensen and Cynthia (2000), 43 - Kohoutek and Pauls (1980), 44 - Leibowitz et al. (1992), 45 - Robinson et al. (1982), 46 - Diaz and Bruch (1994), 47 - Campbell and Shafter (1995)

Additional sources for orbital period, t_2/t_3 time and catalogue masses:

Duerbeck (1987), Warner (1986, 1995, 2002), Diaz and Bruch (1997), Ritter and Kolb (2003)